

# **Guidance for Industry**

## **Characterization and Qualification of Cell Substrates and Other Biological Materials Used in the Production of Viral Vaccines for Infectious Disease Indications**

Additional copies of this guidance are available from the Office of Communication, Outreach and Development (OCOD) (HFM-40), 1401 Rockville Pike, Suite 200N, Rockville, MD 20852-1448, or by calling 1-800-835-4709 or 301-827-1800, or email [ocod@fda.hhs.gov](mailto:ocod@fda.hhs.gov), or from the Internet at <http://www.fda.gov/BiologicsBloodVaccines/GuidanceComplianceRegulatoryInformation/Guidances/default.htm>.

For questions on the content of this guidance, contact OCOD at the phone numbers listed above.

**U.S. Department of Health and Human Services  
Food and Drug Administration  
Center for Biologics Evaluation and Research  
[February 2010]**

## TABLE OF CONTENTS

<b>I.</b>	<b>INTRODUCTION</b> .....	<b>1</b>
<b>II.</b>	<b>DEFINITION</b> .....	<b>2</b>
<b>III.</b>	<b>CHARACTERIZATION AND QUALIFICATION OF CELL SUBSTRATES, VIRAL SEEDS, BIOLOGICAL RAW MATERIALS AND VACCINE INTERMEDIATES</b> .....	<b>2</b>
<b>A.</b>	<b>PRODUCT-SPECIFIC PARAMETERS INFLUENCING CHARACTERIZATION AND QUALIFICATION OF CELL SUBSTRATES</b> .....	<b>3</b>
1.	<i>Vaccine Purity</i> .....	3
2.	<i>Potential Sources of Contamination</i> .....	4
3.	<i>Current Good Manufacturing Practices (cGMP) and Cell Substrate Development</i> .....	4
4.	<i>Use of Control-Cell Cultures</i> .....	4
5.	<i>Assay Validation</i> .....	5
<b>B.</b>	<b>PROPERTIES OF THE CELL SUBSTRATE</b> .....	<b>5</b>
1.	<i>Properties Relevant to Cell Substrate Selection</i> .....	6
2.	<i>Source</i> .....	6
3.	<i>History (including identifying characteristics) and other important characteristics</i> .....	7
4.	<i>Growth Characteristics</i> .....	8
5.	<i>Expression Characteristics</i> .....	8
6.	<i>Susceptibility to Adventitious Agents</i> .....	8
7.	<i>Generation of Cell Substrate</i> .....	8
<b>C.</b>	<b>CELL BANKING</b> .....	<b>9</b>
1.	<i>Cell Banking Strategies and Methods</i> .....	9
2.	<i>Qualification of Cell Banks and Primary Cell Cultures</i> .....	10
3.	<i>Special Considerations for Primary Cell Cultures</i> .....	11
4.	<i>Special Considerations for Diploid Cell Strains</i> .....	12
5.	<i>Special Considerations for Continuous Cell Lines</i> .....	13
6.	<i>Additional Considerations for Cell Lines that are Tumorigenic or Tumor-derived</i> .....	14
<b>D.</b>	<b>VIRAL SEEDS</b> .....	<b>14</b>
1.	<i>Master Viral Seed</i> .....	15
2.	<i>Working Viral Seed</i> .....	16
<b>E.</b>	<b>BIOLOGICAL RAW MATERIALS AND ANCILLARY REAGENTS</b> .....	<b>16</b>
1.	<i>Serum</i> .....	16
2.	<i>Trypsin</i> .....	17
3.	<i>Amino Acids</i> .....	17
4.	<i>Other Biological Reagents</i> .....	17
<b>F.</b>	<b>TESTING AT DIFFERENT STAGES OF PRODUCTION</b> .....	<b>17</b>
1.	<i>Cell Banks</i> .....	17
2.	<i>Pre-Production Cells</i> .....	17
3.	<i>Pre-Filtered Harvest or Post-Production Cells</i> .....	18
4.	<i>Control Cells</i> .....	18
5.	<i>Post-Filtered Harvest (Final Bulk)</i> .....	18
6.	<i>Final Filled Product</i> .....	19
<b>IV.</b>	<b>DESCRIPTION OF TEST METHODS</b> .....	<b>19</b>
<b>A.</b>	<b>TESTING FOR ADVENTITIOUS AGENTS</b> .....	<b>19</b>
1.	<i>In Vivo Tests</i> .....	20
a.	<i>Adult Mice</i> .....	20
b.	<i>Suckling Mice</i> .....	21
c.	<i>Guinea Pigs</i> .....	21
d.	<i>Rabbits</i> .....	22
e.	<i>Embryonated Chicken Eggs</i> .....	22
f.	<i>Antibody Production Tests</i> .....	23
2.	<i>In Vitro Tests for Viruses</i> .....	24

**Contains Nonbinding Recommendations**

- a. Cell Culture Safety Tests ..... 24
  - *Human Diploid Cells*..... 24
  - *Monkey Kidney Cell Line (usually Vero cells)* ..... 24
- b. Transmission Electron Microscopy (TEM) ..... 26
- c. Biochemical Tests for Retroviruses ..... 26
- d. Infectivity Tests for Retroviruses ..... 27
- e. PCR or Other Specific In Vitro Tests ..... 27
- f. Sourcing and Testing of Bovine-Derived Materials ..... 28
- g. Testing of Porcine-Derived Reagents ..... 29
- 3. *In Vitro Tests for Non-Viral Agents*..... 29
  - a. Mycoplasma/Spiroplasma ..... 29
  - b. Bacterial and Fungal Sterility ..... 32
  - c. Mycobacteria Testing ..... 32
- B. TESTING OF CELL PROPERTIES ..... 33
  - 1. *Tests for Tumorigenicity*..... 33
  - 2. *Testing for Oncogenicity* ..... 35
  - 3. *Identity Testing of Cell Substrates*..... 35
  - 4. *Testing for Genetic Stability*..... 36
- C. OTHER TESTS ..... 36
  - 1. *Testing for the Presence of Residual Cells* ..... 36
  - 2. *Testing for Residual Cellular DNA*..... 37
  - 3. *General Safety Test (GST)*..... 37
- V. GLOSSARY ..... 38
- VI. REFERENCE LIST ..... 42

## Guidance for Industry

# Characterization and Qualification of Cell Substrates and Other Biological Materials Used in the Production of Viral Vaccines for Infectious Disease Indications

*This guidance represents the Food and Drug Administration's (FDA's) current thinking on this topic. It does not create or confer any rights for or on any person and does not operate to bind FDA or the public. You can use an alternative approach if the approach satisfies the requirements of the applicable statutes and regulations. If you want to discuss an alternative approach, contact the appropriate FDA staff. If you cannot identify the appropriate FDA staff, call the appropriate number listed on the title page of this guidance.*

## I. INTRODUCTION

We, FDA, are providing you, manufacturers of viral vaccines, guidance for the characterization and qualification of cell substrates, viral seeds, and other biological materials used for the production of viral vaccines for human use.

This guidance applies to the development of viral vaccines for the prevention and treatment of infectious diseases that are regulated by the Office of Vaccines Research and Review (OVR) of the Center for Biologics Evaluation and Research (CBER) under section 351 of the Public Health Service (PHS) Act (42 U.S.C. 262).

Cell substrates are cells used to produce vaccines. The scope of this guidance document is limited to cell substrates of human or animal (including insect) origin and does not cover characterization of unicellular organisms, such as bacteria or yeast. In this document, cell substrates are categorized as primary (cells, including eggs, derived directly from an animal source and that are not stored as cell banks), diploid (cells with a normal or near-normal karyotype and that are stored as cell banks prior to use in vaccine manufacture), or continuous (cells that are immortal and do not undergo senescence). This guidance also applies to the characterization and qualification of viral seeds and other biological materials (including vaccine intermediates) used in vaccine manufacture.

This guidance finalizes the draft guidance entitled "Guidance for Industry: Characterization and Qualification of Cell Substrates and Other Biological Starting Materials Used in the Production of Viral Vaccines for the Prevention and Treatment of Infectious Diseases," dated September 2006 (71 FR 57547). In addition, this document replaces the information pertaining to viral vaccines for the prevention and treatment of infectious diseases that we provided in the document entitled "Points to Consider in the Characterization of Cell Lines Used to Produce

## **Contains Nonbinding Recommendations**

Biologicals,” dated 1993 (Ref. 1). This document also supplements the recommendations on the production of viral vaccines for the prevention and treatment of infectious diseases, provided in International Conference on Harmonization (ICH) documents Q5A and Q5D (Refs. 2 and 3, respectively). For the production of biological products not covered under this guidance, we recommend that you refer to the “Points to Consider in the Characterization of Cell Lines Used to Produce Biologicals,” dated 1993 (Ref. 1).

FDA’s guidance documents, including this guidance, do not establish legally enforceable responsibilities. Instead, guidances describe FDA’s current thinking on a topic and should be viewed only as recommendations unless specific regulatory or statutory requirements are cited. The use of the word *should* in a guidance document means that something is recommended, but not required.

## **II. DEFINITION**

For the purpose of this document, viral vaccines are a heterogeneous class of preventive, and in some cases, therapeutic medicinal products that when administered are intended to elicit immune responses that could prevent and/or lessen the severity of one or more infectious diseases. These products include live attenuated preparations of viruses, inactivated (killed) whole or subunit virions, purified recombinant proteins, synthetic antigens, or live viral vectors expressing specific heterologous vaccine antigens. Therapeutic vaccines for non-infectious diseases (e.g., certain cancer vaccines) and monoclonal antibodies used as immunogens (e.g., anti-idiotypic antibodies) are not considered here. A glossary defining scientific terms in this guidance is located in Section V below.

## **III. CHARACTERIZATION AND QUALIFICATION OF CELL SUBSTRATES, VIRAL SEEDS, BIOLOGICAL RAW MATERIALS AND VACCINE INTERMEDIATES**

Selection of a cell substrate influences the safety and purity of the biological product manufactured in it. The evaluation (e.g., the particular tests and methodologies chosen) should be specifically tailored to the cell substrate and vaccine product to be developed to assure vaccine safety and purity. FDA recognizes that technology in this area is evolving. This section describes general parameters to be considered in the characterization of cell substrates, viral seeds, biological starting materials and vaccine intermediates. To maximize the value of the recommendations in this document, it is important that you perform each test at the stage of production at which contamination is most likely to be detected.

Tests to characterize and qualify cell substrates, biological materials, vaccine seeds, and vaccine intermediates are described in Section IV.

## Contains Nonbinding Recommendations

### A. PRODUCT-SPECIFIC PARAMETERS INFLUENCING CHARACTERIZATION AND QUALIFICATION OF CELL SUBSTRATES

#### 1. Vaccine Purity

The regulations, in 21 CFR 610.13, state in part that “Products shall be free of extraneous material except that which is unavoidable in the manufacturing process described in the approved biologics license application.” In 21 CFR 600.3(r), purity is defined as the “relative freedom from extraneous matter in the finished product, whether or not harmful to the recipient or deleterious to the product.”

Live attenuated viruses, whole inactivated virions, or virus-like particles often cannot be purified as rigorously as viral subunit vaccines; as a consequence, the potential for contamination is greater than that of subunit vaccines. Generation of live viral vaccines often involves cell disruption, which may add cellular components to the vaccine bulk. In addition, such vaccines often are minimally purified and are not subjected to inactivation steps. Comprehensive testing and qualification of the biological starting materials and lot-by-lot testing for adventitious agents are particularly important because it is not possible for a manufacturer of live viral vaccines to validate the process for clearance of adventitious agents.

In contrast, for more highly purified products where viral clearance can be demonstrated during downstream processing, you can place more reliance on process validation (Ref. 2). For inactivated vaccines, the concern is that the process used to inactivate the vaccine virus may not inactivate all adventitious agents potentially present (as occurred with early inactivated poliovirus vaccines [Ref. 4]). Therefore, you should validate your process for inactivation of adventitious agents using different model viruses (Ref. 2). The choice of tests and the stages at which the tests are applied will depend on your inactivation process. The degree of viral clearance that is feasible may influence the sensitivity of the testing you should perform to demonstrate the absence of contaminants in your product.

If a viral vaccine seed was exposed to a known adventitious agent or if the passage history of a virus seed is unclear, you should purify the viral seed (e.g., by molecular cloning, serial passage in medium containing neutralizing antibody directed against the adventitious agent, or plaque purification). If your purification method is demonstrated to be capable of removing all adventitious agents from a viral seed to within an acceptable safety margin, this approach could be used to qualify a seed. If you are planning to perform seed purification, you should discuss your proposed methodology and its validation with us as early as is feasible.

## **Contains Nonbinding Recommendations**

### **2. Potential Sources of Contamination**

It is important that you identify and examine all potential sources of contamination of your product. For example, a viral seed could be exposed to the following potential sources of contamination: the person or animal from which it was isolated; the cells and raw materials (e.g., serum or trypsin) used in its isolation and attenuation; materials used in banking and propagation of cells for viral seed growth; and other materials used during production and filling of the seed. You should also consider the species of origin of your cell substrates, viral seeds, and other biological starting materials in selecting your tests to ensure the absence of contaminants. Furthermore, you should consider any infectious viruses (including those that infect nonhuman species) as potential contaminants if there is the possibility of contact with your product or cell substrate at any time during development or production. Retroviruses may be either endogeneous (i.e., encoded by the cell substrate genome) or exogenously acquired. Retrovirus testing should address the possibility that either type of retrovirus could contaminate a product. Finally, you should consider the possibility of contamination from unusual sources, as exemplified by the reported presence of minute virus of mice (MVM) in some lots of recombinant proteins (Ref. 5). The susceptibility of the cell substrate to infection by agents of potential concern can influence the tests needed to assure absence of contamination.

Use of qualified raw materials can reduce the risk of introducing adventitious agents. For example, inactivation of viruses by irradiation of serum could provide additional assurance regarding the purity of the final product.

### **3. Current Good Manufacturing Practices (cGMP) and Cell Substrate Development**

Concepts of quality design consistent with cGMP (see 21 CFR Part 211) are relevant to the selection of suitable cell substrates and viral seeds. If your cell substrate or viral seed was not originally derived under concepts consistent with cGMP, additional documentation (e.g., donor and source of raw materials), testing, and production steps might be necessary to support its use. This may include having some virus seeds re-derived or further purified to reduce the possibility of adventitious agent contamination.

### **4. Use of Control-Cell Cultures**

We recommend the use of control cells when it is not feasible to directly test cells or product at various stages of manufacture. For example, if you are using primary cell cultures to propagate your vaccine virus, complete testing of the primary culture might not be feasible prior to inoculation of virus. In this situation, when possible, you should produce and test uninfected control-cell cultures that are derived in parallel with, and handled in the same manner as, the production culture.

## Contains Nonbinding Recommendations

Control-cell cultures are also useful for testing in situations when your vaccine virus can interfere with the results of testing of the virus seed or vaccine product (e.g., when the virus cannot be adequately neutralized to permit testing for adventitious agents).

Control cell cultures should be propagated under conditions similar to conditions of manufacture for a time appropriate to the reason for performing the cultures. This time period may be 14 days or more, when needed to allow for detection of potential adventitious agents that may be latent, endogenous, or poorly replicating. Control-cell cultures should be processed simultaneously with your production culture, but left uninfected. Control cells should be evaluated for the presence of adventitious agents using the same tests that would have been performed on the production cultures, if it were feasible. Tests for adventitious agents should be performed on control cells at times at which you would perform similar tests on your manufactured product.

Testing of control cells does not always eliminate the need for testing final product (see Section III.F.6).

### 5. Assay Validation

You should demonstrate the reliability of assays or tests used to evaluate your cell substrate in the context of intended use. Assays related to assurance of safety should be scientifically valid (for example, by formal validation and/or inclusion of appropriate controls or standards) prior to initiation of clinical trials. Guidance regarding validation of analytical assays may be obtained from the ICH Q2(R1) documents (Ref. 6). Test methods should be validated according to the principles defined in these guidance documents. Certain “compendial” tests (for example, those promulgated by the USP) might not require full validation, as long as you verify their suitability for their intended purpose under your actual conditions of use (Ref. 7).

You should use appropriate statistical analyses to support your assay validation. This may include assessment of assay accuracy, precision, limits of detection, limits of quantification, specificity, linearity and range, ruggedness and robustness, and system suitability. Further guidance in this regard may be obtained from the ICH Q2(R1) document (Ref. 6).

## **B. PROPERTIES OF THE CELL SUBSTRATE**

The cell substrate you choose for production of your viral vaccine should be well characterized. Your application should address the following issues.



## Contains Nonbinding Recommendations

### 1. Properties Relevant to Cell Substrate Selection

Assuring that the vaccine virus strain remains stable during passage in cell culture is an important component of cell substrate selection. Different cell lines may apply different selective pressures on the vaccine virus, which could alter its sequence and possibly its phenotype. For example, when Sabin poliovirus strains are grown in different cells, their likelihood of reversion to neurovirulence is different (Ref. 8). If attenuating mutations or other genetic markers (including expression of antigens relevant to immune response) of the vaccine virus are known, data regarding the influence of serial passage in the cell substrate on retention of these markers can be useful in characterizing the genetic stability of the vaccine virus.

Whatever starting materials are used for the generation of the cell substrate (e.g., parent cells or plasmids used for genetically engineered cells), any available information about those starting materials and their characterization (e.g., sequence of the plasmid) should be provided. If a sponsor starts with a primary cell to generate a novel cell substrate, complete information on donor screening and testing should be provided.

### 2. Source

You should document the species of origin and the tissue type of your cell substrate. You should also document the medical history of the donor and the results of any screening and testing performed on the donor or on samples from the donor. When feasible, human donors should be screened and tested in a manner comparable with whole blood or source plasma donors, as described in 21 CFR Part 640. Animal donors should be screened and tested by applying similar concepts, and should be in general good health (except in cases where the cell substrate is derived from a tumor) and free from symptoms of infectious diseases. Ideally, animals intended to be donors should be quarantined prior to sacrifice or obtaining the sample to be used to generate the cell substrate (Ref. 9). (Issues specific to bovine- and porcine-derived materials are discussed in Sections IV.A.2.f and IV.A.2.g.) The species of origin is highly relevant, because different animal species have different potential contaminants, and some animal cells might be capable of propagating human viruses that are of potential concern as adventitious agents. Tissue type is also an important element for you to describe and consider in deciding on testing strategies. For example, neuronal cells might be more likely to harbor latent viruses (e.g., herpesviruses) or express infectious prion proteins (PrP) and should be evaluated for these potential adventitious agents. Additional considerations might apply to cells that are tumor-derived.

## Contains Nonbinding Recommendations

### 3. History (including identifying characteristics) and other important characteristics

The history of your vaccine cell substrate must be identified under 21 CFR 610.18(c)(1)(i) and should include, to the extent available: sources and testing of the raw materials to which it was exposed in its passage history; listing of any other agents handled or grown in the same biosafety cabinet or incubators around the time of cell substrate passage; and the conditions under which it was passaged (including adherence to cGMPs, if applicable). In some cases (e.g., Chinese hamster ovary [CHO] cells and Vero cells), multiple related cell lines or cell clones exist, and it is important to specify which of these related lines are being used, including the source from which it was obtained and its passage (or population doubling) level. All available documentation of all raw materials of human or animal origin you used for the entire passage history should be retained and considered in developing a testing program. This is particularly important for passages using bovine materials acquired during and after 1980, when bovine spongiform encephalopathy (BSE) emerged in the U.K. You should document all known manipulations (adaptation, engineering, cloning) performed on the cell substrate since original isolation from a primary donor, whether they occurred prior to or after your receipt of the cells.

In addition, you should provide the following information for any cell substrates used for the production of viral vaccines:

- results of all identity testing, including information about any characteristics (such as cytogenetic characteristics, required by 21 CFR 610.18(c)(1)(ii)) that could be used to identify the cells;
- results of all available adventitious agent testing (including that required by 21 CFR 610.18(c)(1)(iv));
- age, sex, and species of the donor;
- donor's medical history and the results of tests performed on the donor for the detection of adventitious agents;
- culture history of the cell line, including methods used for the isolation of the tissues from which the line was derived, passage history, medium used, and history of passage in animals;
- documentation of the history of human-derived and animal-derived materials used during passage of the cells; and
- documentation of any genetic material introduced into the cell substrate.

For situations when the specified information is not available, data derived from testing of the cell substrate by other methods may prove supportive and may be sufficient.

## Contains Nonbinding Recommendations

### 4. Growth Characteristics

You must characterize the in vitro growth characteristics (e.g., suspension or monolayer cultures, rate of growth) of your cell substrate (21 CFR 610.18(c)(1)(iii)), except if you use primary cell cultures exempted from that requirement by 21 CFR 610.18(c)(3). If the cells have a finite life expectancy, the total number of population doubling levels through senescence should be determined.

A description of the tumorigenic property of cells is required for all diploid and non-diploid cells, including continuous cell lines (21 CFR 610.18(c)(1)(ii)). Because previous experience has consistently demonstrated the non-tumorigenic phenotype of the well-characterized diploid cell strains MRC-5, WI-38, and FRhL-2 (if they are not genetically or phenotypically modified), further tumorigenicity testing of banks consisting of these cell strains is not considered necessary to satisfy this requirement (please also see Section III.C.4). A description of the tumorigenic property is not required for primary cell cultures that are not subcultivated or that are subsequently subcultivated for only a very limited number of population doublings (21 CFR 610.18(c)(3)).

If there are specific markers that might be useful in characterizing your cell line (such as microscopic appearance, marker chromosomes, transfected genetic elements, specific surface markers, etc.), these should be analyzed for their stability through manufacturing passage levels.

### 5. Expression Characteristics

If your cell substrate is used to produce a recombinant protein, you should evaluate its ability to do so as a part of its characterization. This includes evaluating the copy number and stability of introduced nucleic acids and the quantity and quality of expressed proteins.

### 6. Susceptibility to Adventitious Agents

In developing comprehensive and appropriate testing strategies for adventitious agents, you should consider the susceptibility of the cell substrate to infection with viruses other than the vaccine strain. Selection of tests should be based on potential sources of contamination (See Section II.A.2). Your production scheme and testing program should be designed to assure the absence of viruses that might replicate in your cell substrate.

### 7. Generation of Cell Substrate

You should characterize the cell substrate for the impact that the manipulations have had on the characteristics of the cell when you use cell substrates that have been generated by in vitro manipulations or engineered to express a viral antigen.

## **Contains Nonbinding Recommendations**

For example, immortalization of primary cells with viral oncogenes to generate a complementing cell line for production of defective viruses or viral vectors will inevitably change the phenotype of the cell. Such manipulations might induce the expression of latent or endogenous viruses, and/or might cause the cells to become tumorigenic. In addition, a cell substrate that has been derived by cell cloning might have different characteristics from the parental cell line. Because it is derived from one or a few cells, it might not have characteristics representative of the original population from which it was cloned. Alternatively, a clone might be selected as the cell substrate because of its particular outlier characteristics, such as rapid propagation in culture or adaptation to particular cell culture conditions that modify its growth properties to enhance vaccine virus replication (e.g., development of suspension cell cultures from adherent cells). It is important that you thoroughly evaluate the characteristics of derivative or engineered cell substrates, as it cannot be assumed that the parental cell characteristics were maintained following the manipulations used to generate the production cell substrate.

### **C. CELL BANKING**

Cell banking assures that an adequate supply of equivalent, well-characterized cells exist for production over the expected lifetime of the product. In addition to providing a constant supply of biological starting material, cell banking provides you with the opportunity to undertake a comprehensive characterization of the cell substrate and to minimize the chance of adventitious agent contamination and/or to maximize the chance of detection of a contaminant.

#### **1. Cell Banking Strategies and Methods**

Ordinarily, the cell bank system consists of two tiers: a master cell bank (MCB); and a working cell bank (WCB), sometimes called a manufacturer's working cell bank (MWCB). The MCB represents a collection of cells of uniform composition derived from a single source prepared under defined culture conditions. Sometimes, a parent cell bank, comprising vials of the progenitor cells to the MCB, is also maintained.

The WCB is derived from one or more vials of cells from the MCB, which are expanded by serial subculture. The pooled cells are dispensed into individual vials and cryopreserved to form the WCB. One or more vials from the WCB are generally used for the production of a lot of a vaccine. If cells from more than one WCB vial are used, the cell suspensions should be pooled at the time of thawing. The population doubling level or passage level of cells used for production should not exceed an upper limit based on your cell substrate characterization, including at end-of-production level (See Section III.C.2).

When using a two-tiered banking system, we generally recommend that you perform an extensive characterization of the MCB. Because all WCBs would be

## **Contains Nonbinding Recommendations**

derived from the well-characterized MCB, your WCBs may be tested in a more limited manner, focusing on adventitious agents to which the WCB could have been exposed during expansion from the MCB. The use of a cell banking system might allow you to perform more limited testing of individual vaccine lots, focusing on adventitious agents to which the cells could have been exposed during manufacturing from the WCB. Alternatively, some manufacturers might choose to extensively characterize each WCB in lieu of a one-time thorough characterization of the MCB. The testing strategy chosen will depend on your own circumstances, but you should characterize at least one cell bank (MCB or WCB) extensively. However, in the absence of thorough characterization of the MCB, we may request a thorough characterization on the WCB used to produce each lot.

You may use other cell banking systems, for example, when the original source of your cell substrate is limited and/or the need to maintain a low passage level restricts expansion of cell numbers. Regardless of what cell banking system you use, cells at some level of expansion should be characterized completely, with a justification for the choice of the passage level tested. In either case, the derivation and cell bank designations should be thoroughly described and explained.

The requirements set out in 21 CFR 610.18 relate to cultures, including cell banks. Cell banks should be stored under conditions shown to be suitable for long-term stability (e.g., liquid nitrogen, ultra-low temperature freezer) (21 CFR 610.18(a)). Storage of cell banks in the vapor phase (as compared to the liquid phase) of liquid nitrogen might reduce the potential for cross-contamination. Cell stability under the freezing and storage conditions should be validated using cell recovery or viability data. You should store the MCB and WCB in two or more separate locations within the facility or at a distant site in order to avoid loss of the cell substrate due to local disaster or equipment malfunction. Access to these cell banks by your staff should be limited and controlled. Under 21 CFR 610.18(d), appropriate records must be maintained. You should maintain a record of the location, identity, and inventory of individual ampoules of cells.

### **2. Qualification of Cell Banks and Primary Cell Cultures**

The purpose of qualifying cell banks is to demonstrate their suitability for use in vaccine production. This section includes recommendations on testing for different types of cells and stages of cell banking.

The same basic principles apply to the qualification of cell banks and to the qualification of primary cell cultures. Characterization of a cell bank is dependent on its use, and your particular use should guide your determination of which recommended testing is necessary and whether additional testing might be necessary. Consideration of the choice of tests will be based on the assessment of the risks that the cell substrate might represent for the product type and its clinical indications.

## Contains Nonbinding Recommendations

Testing to qualify the MCB should be performed directly on the cell bank, except when it is more appropriate to test cell cultures derived from the cell bank.

Tests that you might perform on your MCB include tests for bacteria, fungi, mycoplasma, and viruses (e.g., in vitro and in vivo testing, specific tests for retroviruses, and specific tests for viruses known to exist in the species of origin or that could be acquired during serial passage in cell culture). Other specific tests might be warranted, depending on the passage history of the cell substrate. Details of these tests are presented in Section IV. If the species from which your cells were derived is susceptible to infection with *Mycobacterium tuberculosis*, an appropriate test should be performed for this agent as well. Depending on the source species, you may consider specific tests such as the test for avian leukosis virus for products produced in avian cells. If you are using primary monkey kidney cell cultures, for example, you should test for species-specific simian agents, such as SV40 and herpes B virus, and other simian agents, such as simian polyomaviruses and simian cytomegalovirus.

Testing either the MCB or the reagents to which it was exposed may be done to provide assurance that these reagents were not a source of adventitious agents, an issue of particular concern when reagents are animal-derived. Some of the tests that are relevant in selected circumstances are described in Section IV, including the tests for bovine (Section IV.A.2.f) and porcine (Section IV.A.2.g) agents. Additional considerations are described for serum in Section III.E.1 and for trypsin in Section III.E.2. In the case of bovine derived materials, the issue of TSE agents should be considered (Ref. 33). Finally, testing should be performed in most circumstances to determine if the cells produce retroviruses or retrovirus particles. This testing is described in Sections IV.A.2.c and IV.A.2.d.

Extensive characterization of the MCB should be done once and should also address the considerations in Section III.B. Methods that may be used for identity testing of the MCB are discussed in Section IV.B.3.

You should characterize cells that are expanded to or beyond the end of production passage level. Such cells are referred to as End-of-Production Cells (EOPC). You should demonstrate the stability of the cell substrate characteristics using your EOPC. Your characterization should include, if applicable, growth characteristics, tumorigenic phenotype, expression of endogenous viruses, stability of expression of the inserted or engineered genes, and genetic stability of the cells (such as karyotyping or DNA fingerprinting).

### 3. Special Considerations for Primary Cell Cultures

Primary cells are obtained directly from the tissues of healthy animals. Primary cells are more likely to contain adventitious agents than banked, well-characterized cells. This concern with primary cells is mitigated by rigorous qualification of source animals and primary cell substrates. Animals from which primary cultures

## Contains Nonbinding Recommendations

are established should be from specific-pathogen-free (SPF) closed flocks, herds, or colonies, when feasible. The term “closed” refers to the maintenance of a group (flock, herd, and colony) free from introduction of new animals (new genetic material). Animals that are not from closed flocks, herds, or colonies should be quarantined and thoroughly evaluated for a period sufficient to detect signs of disease or infection before introducing them into the flock, herd, or colony. Animals should be screened serologically for appropriate adventitious agents to determine their suitability as a source for the primary cell substrate. Animal husbandry practices should be described in the application.

Embryonated chicken eggs used as the source of chick embryo tissue for the propagation of viral vaccines should be derived from flocks certified to be free of *Salmonella pullorum*, avian tuberculosis, fowl pox, Rous sarcoma virus, avian leukosis virus, reticuloendotheliosis virus, and other adventitious agents pathogenic for chickens. Appropriate records must be retained for documentation (21 CFR 610.18(d)). If eggs are procured from non-SPF flocks, assurance that the vaccine is free of such agents may be provided by testing or by validating the process for their removal.

Control cells are often used with primary cell substrates and are described in more detail in Section III.A.4.

#### 4. Special Considerations for Diploid Cell Strains

Diploid cell strains are established from primary cell cultures by expansion and cell banking. These types of cells have a finite life span and are not immortal like cell lines. Diploid cells usually retain a diploid or near diploid karyotype, a characteristic that also differs from cell lines, which are generally aneuploid or non-diploid.

You should determine the karyotype of new diploid cell strains to establish identity and to characterize a cell strain. Such analyses will establish the diploid character of your cells and determine its freedom from contamination with other cell lines. We recommend that you monitor the genetic stability of the diploid cell strain throughout production. Other tests of identity (see Section IV.B.3) may also be appropriate.

Animal tumorigenicity testing is not needed if you are using genetically or phenotypically unmodified previously characterized diploid cell strains, such as MRC-5, WI-38, and FRhL-2 (see Section III.B.4). However, tumorigenicity testing must be performed for a novel diploid cell substrate because such testing is needed to satisfy the requirement in 21 CFR 610.18(c)(1)(ii) that you describe its tumorigenicity.

## Contains Nonbinding Recommendations

### 5. Special Considerations for Continuous Cell Lines

Some continuous cell lines, including Vero cells and CHO cells, have been used as substrates for licensed biologicals. Cell lines might have biochemical, biological, and genetic characteristics that differ from primary or diploid cells (e.g., they are typically aneuploid and have accumulated genetic changes). Because the mechanism by which most cell lines become immortal is generally unknown, and because some cell lines form tumors when inoculated into immunodeficient rodents, there are additional concerns for continuous cell lines compared with diploid cells, including the potential that transformation was caused by an adventitious agent and potential risks from residual DNA.

Products prepared in cell lines (including viral vaccines) should be purified to be free (see Section V. Glossary) of adventitious agents and residual cells and should have low levels of cell-substrate DNA. When potential biological activity of residual cell substrate DNA is a concern, you should introduce procedures that extensively remove or degrade DNA. If you are considering the use of cell lines, we encourage you to develop and evaluate efficient methods for the purification of your product as an essential element of any product development program.

Because certain cell lines express endogenous viruses (e.g., retroviruses), tests capable of detecting such agents should be completed on cells grown under production conditions (See Section IV.A. Testing for Adventitious Agents). With the exception of chicken cells (see Section IV.A.2.c), if specific contaminants have been identified as endogenous agents in the MCBs and WCBs you are using, you should demonstrate their clearance (inactivation and/or removal) to a defined level by the purification procedures you use in production. You may obtain further guidance on this topic from the ICH Q5A document (Ref. 2).

The testing recommended for qualification of cell substrates and cell banks should also be applied to rodent cell lines. Rodent cell lines are presumed to be capable of producing endogenous retroviruses. Assessment of the quantity and type of retroviruses produced should be performed. Infectivity assays for retroviruses are also recommended. You should only use rodent cell lines if your product can be sufficiently purified to demonstrate levels of viral clearance that assure the final product is not contaminated with retroviral particles. You may obtain additional guidance on viral clearance validation from the ICH Q5A document (Ref. 2).

Under 21 CFR 610.18(c)(1)(ii), you must describe cell lines with respect to tumorigenicity. Cell lines could acquire tumorigenic properties with increasing passage levels. It is therefore important that you limit the passage level of the cells used in production and that you characterize cells at or beyond this end-of-production limit. The maximum end-of-production passage level should be based on data derived from production cells expanded under comparable or analogous conditions to the production conditions. Cells from either the MCB or the WCB may be expanded for this evaluation. Tumorigenicity testing should be performed at or beyond this



## Contains Nonbinding Recommendations

level. If you increase your defined limit for end-of-production, you should support the safety of this with data from cells that have been expanded to a passage level or population doubling level that is equal to or greater than your newly proposed end-of-production level.

### 6. Additional Considerations for Cell Lines that are Tumorigenic or Tumor-derived

Use of tumorigenic and tumor-derived cells for vaccine production is associated with additional issues. This document does not provide guidance on a pathway for licensure of live-attenuated or minimally purified vaccine produced in these cells. You should perform additional testing if your cell lines are tumorigenic or derived from a tumor. You should assess cell lines that are tumorigenic or tumor-derived for potential oncogenic viruses and oncogenic substances (including nucleic acids), which could be associated with induction of a neoplastic process in a vaccine recipient. Test strategies for potential oncogenic viruses or oncogenic substances may be determined case-by-case, depending on the tissue type, source species, passage history, and extent of knowledge about the transforming event(s). (See section IV.B.1 for a discussion of tumorigenicity testing and section IV.B.2 for a discussion of oncogenicity testing.)

In cases where you know the transforming event (e.g., if the cells were transformed by a known oncogene), your testing should demonstrate that your final product is free of the transforming agent. For example, if adenovirus sequences are used to transform a primary human cell to produce a cell line (e.g., 293 cells), then testing should demonstrate that the final product is free of the introduced viral sequences. Similarly, if a virus is used to transform cells, that virus and its genetic material should not be detectable in the final product using an assay with sensitivity sufficient to provide assurance of safety. Tumorigenic or tumor-derived cell lines for which the mechanism of transformation is unknown will require additional testing to ensure the absence of potential transforming and oncogenic agents. You should consult with CBER on which tests you should perform and which methods you should follow.

## D. VIRAL SEEDS

As with cell banks, you should document passage history and derivation history of viral seeds. Your description should include donor screening, testing and donor medical history. Any manipulation of the viral phenotype, such as cold-adaptation, development of temperature-sensitivity, or attenuation of virulence, should be well documented. Any genetic manipulations, such as reassortment or recombination, should also be well documented, including determining the nucleic acid sequences and sourcing of each biological starting material (e.g., plasmids, parental viruses).

Vaccine virus banks are commonly referred to as master viral seed (MVS) and working viral seed (WVS). Viral seeds should be stored under conditions that maintain their

## Contains Nonbinding Recommendations

stability (i.e., in liquid nitrogen or ultra-low temperature freezers) and in more than one location within a facility or at a distant site for security reasons. You should assess your viral seed for its growth characteristics on your production cell substrate, tissue tropism, genetic markers, viability during storage, genetic stability through production, attenuation (if applicable), and absence of adventitious agents. If attenuation or derivation is achieved by passage through different cell types from different species, your viral seed should be assessed for absence of adventitious agents from all species that it might have been exposed to from isolation, through passage, and during production, including those that might be present in the raw materials used at each of these stages.

### 1. Master Viral Seed

We recommend that you extensively characterize your MVS, as this characterization also provides assurance regarding the characteristics of subsequent passages, including the WVS. In addition, you should demonstrate the stability of the genotype and phenotype following viral passages beyond the level used in your production. Genotypic characterization of a viral seed includes its sequence, and may include analysis of viral subpopulations and its genetic stability, including susceptibility to reversion. Phenotypic characterization of a viral seed may include assessment of tissue tropism, attenuation properties, and temperature sensitivity, if applicable.

Under 21 CFR Part 610, you must perform tests for safety, purity, and potency of a product, as required. This includes tests for identity, bacterial and fungal sterility, the presence of mycoplasmas, *Mycobacterium tuberculosis* (if appropriate), and adventitious viruses (in vitro and in vivo tests). You should also consider specific tests for agents that might be present in the seed due to its passage history.

In some cases, it may be necessary to neutralize the vaccine virus in order to adequately test for adventitious agents in vitro and in vivo. Preferably, neutralizing antibodies should be monoclonal to reduce the likelihood of also neutralizing adventitious viruses that could be present in the seed. In addition, due to the potential for cross-neutralization of adventitious human viruses, neutralizing antibodies should generally not be prepared from human or primate serum. It is important that you demonstrate that the neutralization procedure does not interfere with detection of adventitious viruses. Sometimes, it is not possible to efficiently neutralize a viral seed. In such cases, you may choose alternative strategies, including testing smaller quantities of seed, subculture onto fresh target cells in the in vitro adventitious agent test, introduction of additional tests (e.g., polymerase chain reaction (PCR), Antibody Production assays (Ref. 10) or testing of control cells (Section III.A.4).

Assessment of neurovirulence might be appropriate for MVSs derived from wild type viruses known or suspected to possess neurotropic or neurovirulent properties, and we recommend that you consult with CBER on appropriate animal models, methods, and scoring systems for this assessment before you initiate such studies.

## **Contains Nonbinding Recommendations**

Animal models other than non-human primates may be acceptable. For live-attenuated viruses that are neurovirulent or might revert to neurovirulence (e.g., polioviruses), it might be necessary to assess neurovirulence not only on the MVS or an end-of-production passage level virus stock, but also on the product lot-by-lot.

### **2. Working Viral Seed**

The level of characterization of a WVS should be based on the extent of characterization of the MVS from which it was derived. Once you have demonstrated your MVS to be free of adventitious agents from the species to which your vaccine virus had been exposed during its isolation and passage history, you only need to show that your WVS is free of adventitious agents that could have been introduced in generating the WVS (e.g., from production cells and raw materials used in propagation and processing). Alternatively, more extensive testing could be performed on WVS material if less extensive testing has been performed on the MVS in an effort to conserve the supply of MVS.

## **E. BIOLOGICAL RAW MATERIALS AND ANCILLARY REAGENTS**

You should test your raw materials and ancillary reagents to assure that they are appropriate for use in all stages of vaccine production, including the production of your cell banks and viral seeds. Raw materials derived from animals or humans should be shown to be free from adventitious agents of the species from which they are derived. Appropriate records must be retained for all reagents and biological raw materials used for vaccine production (21 CFR 610.18(d)).

If your documentation for the raw materials used in the passage of your cell substrate is insufficient to assure that the raw materials did not introduce adventitious agents, you should test the cell substrate itself for adventitious agents of the relevant species. For human-derived raw materials and reagents, you should be able to document sourcing from appropriately screened and tested donors (using 21 CFR 640.3 and 640.63 as guidelines) or use of products that are already licensed for human use. All biological raw materials should be free of adventitious agents, including bacterial and fungal agents, cultivatable and non-cultivatable mycoplasmas, mycobacteria, and viruses. If your process includes steps that you rely upon to remove or inactivate potential infectious contaminants from biological raw materials, you should validate those steps.

### **1. Serum**

The serum used in vaccine production (including the production of the cell banks and viral seeds), such as bovine serum used in culture medium or human serum albumin used to stabilize vaccine virus, should be tested and certified. Human serum albumin should be licensed for human use in the U.S. or derived from donors

## Contains Nonbinding Recommendations

who have been appropriately screened and tested (21 CFR 640.3 or 640.63). Bovine serum should be free of adventitious agents, including bacterial and fungal agents, mycoplasmas, mycobacteria, and bovine viruses (Section IV.A.2.f).

### 2. Trypsin

You should clearly identify the species from which the trypsin used in your vaccine production (including the production of the cell banks and viral seeds) is derived. If bovine trypsin is used, the concerns identified in Section IV.A.2.f, “Sourcing and Testing of Bovine-Derived Materials,” apply. If porcine trypsin is used, it should be tested in accordance with the recommendations described in Section IV.A.2.g, “Testing of Porcine-Derived Reagents.”

### 3. Amino Acids

You should document the source of the amino acids used in growth medium or in production.

### 4. Other Biological Reagents

You should assess (including testing, where appropriate) the potential risk for introduction of adventitious agents via other biological reagents used during manufacturing, including transferrin, insulin, or other growth factors used in growth medium, as appropriate for their species of origin. If derived from humans, the ancillary reagent should either be licensed in the U.S. or derived from donors for whom appropriate screening and testing have been applied (if appropriate, comparable with blood or source plasma donors, see 21 CFR 640.3 or 640.63). Under 21 CFR 610.15(c), you may not add penicillin to the production substrate of viral vaccines. Also, other beta-lactam antibiotics should not be present in production cell cultures.

## **F. TESTING AT DIFFERENT STAGES OF PRODUCTION**

See Appendix 1 for a list of tests to consider at each stage of production for a viral vaccine.

### 1. Cell Banks

Qualification of cell banks is discussed in Section III.C.

### 2. Pre-Production Cells

Visual inspection or an identity test may be performed on cells directly prior to production.

## Contains Nonbinding Recommendations

### 3. Pre-Filtered Harvest or Post-Production Cells

In general, the stage at which adventitious agents are most likely to be found is the stage at which the most extensive adventitious agent testing should be performed for each product. For many viral vaccines, the pre-filtered harvest is the stage of manufacture that is most concentrated and at which the least processing has been performed. For these reasons, this might be the best stage of production for testing for adventitious agents.

In addition to testing the viral or vaccine harvest for cultivatable mycoplasmas, as required under 21 CFR 610.30, you should also test the viral or vaccine harvest for non-cultivable mycoplasmas (and spiroplasmas, if appropriate), and adventitious viruses by in vitro methods. Adventitious virus tests should also include in vivo methods if the cells and vaccine seeds have not already been tested. This might include a test for hemadsorbing viruses. If the production process is capable of inducing expression of retroviruses, PCR-based reverse transcriptase (RT) testing or infectivity studies might also be recommended at this stage. If appropriate, a test for *Mycobacterium tuberculosis* should be performed.

As discussed for the MVS (Section III.D.1), if the assay system used for in vitro or in vivo adventitious virus testing is capable of supporting replication of the vaccine virus, and if the vaccine virus replicates to levels that interfere with the adventitious agent tests, then it might be necessary to neutralize the vaccine virus prior to performing these tests. If neutralization is difficult to achieve, the other considerations in Section III.D.1 might be applicable.

If cells survive the production process (e.g., if the vaccine virus does not result in a lytic infection), post-production cells may also be tested for adventitious agents. If multiple harvests are performed for a single vaccine lot, testing should be performed on each individual harvest (rather than the pooled harvest) in order to avoid dilution of a potentially contaminated harvest with uncontaminated harvests

### 4. Control Cells

Recommendations for testing of control cells are discussed in Section III.A. 4.

### 5. Post-Filtered Harvest (Final Bulk)

The post-filtered harvest (generally referred to as final bulk) should be tested for bacterial and fungal sterility. Depending on the product, other testing might be appropriate to assure the safety, identity, purity, potency, and quality of the final bulk. These may include testing for levels of residual cellular proteins and cellular nucleic acids. If the cell line used to produce the vaccine is known to produce viruses other than the vaccine virus, the clearance of these viruses should be demonstrated in the production processes. Guidance regarding viral clearance may be sought from the ICH Q5A document (see Ref. 2). If processes are not validated

## **Contains Nonbinding Recommendations**

for removal of residual cells, a test to demonstrate the absence of residual cells is recommended at this stage.

### **6. Final Filled Product**

The regulations for General Biological Products Standards under 21 CFR Part 610 provide that each lot must be tested for potency, general safety, sterility, purity and identity.

Sterility must be demonstrated as required under 21 CFR 610.12. Products excepted from these requirements (e.g., under 21 CFR 610.12(g)(4)(ii)) should still be assessed by the microbial limits test for bioburden (Ref. 11). Under 21 CFR 610.11a, general safety testing for each lot of inactivated influenza vaccine includes a demonstration that it contains less endotoxin than a reference preparation provided by FDA. Other final filled products should also be assessed for endotoxin levels. In addition, a general safety test for detecting extraneous toxic contaminants, under 21 CFR 610.11, is required on final filled containers, except as described in 21 CFR 610.11(g). Other tests should be performed to assess the safety, identity, purity, potency, and quality of the final filled product. These tests are determined case-by-case depending on the product and should be described in your IND and BLA submissions.

## **IV. DESCRIPTION OF TEST METHODS**

The tests described in this Section may be used to address the considerations discussed in Section III. Advances in science and technology are likely to yield additional information that could lead to modification or replacement of some of these tests, and except where prohibited by regulation, manufacturers may use alternative test methods, where scientifically justified. Manufacturers should consider how they can replace, refine, or reduce their use of in vivo tests.

### **A. TESTING FOR ADVENTITIOUS AGENTS**

Assurance that products are free of adventitious agents is a critical component of meeting the 21 CFR 610.13 requirement for purity. Cultures must be tested for the presence of detectable microbial agents and tests necessary to assure the safety and purity of the product may be required (21 CFR 610.18). Your biological starting materials should be characterized sufficiently to ensure that they do not contaminate the final product with extraneous infectious organisms, such as bacteria, fungi, cultivatable and non-cultivatable mycoplasmas and spiroplasma, mycobacteria, viruses, and the agent(s) responsible for transmissible spongiform encephalopathies (TSEs). For a substance to be considered free of a contaminant, your assay should demonstrate, at a predefined level of sensitivity, that a certain quantity of the substance is free of that contaminant. Alternatively, a validated process that is known to remove a contaminant to a defined level may be used to demonstrate the absence of that contaminant. If an adventitious agent is known to be present in your cell substrate or viral seed, then you should demonstrate that your

## Contains Nonbinding Recommendations

production process is sufficiently robust to eliminate or inactivate the agent with an appropriate margin of safety. You should avoid exposure of your product to agents that are known to be infectious for humans (other than the vaccine virus) and/or to agents for which there are no appropriately sensitive validated testing procedures (such as TSEs). You should have appropriate and sufficient controls in place to assure that such exposures do not occur.

This section applies to testing of cell substrates, control cell cultures, viral seeds, and vaccine harvests. When cell substrates are tested, generally, a preparation at  $10^7$  cells/mL (or when appropriate, cell lysate from an equivalent number of cells in culture medium) is used, with the indicated volumes. When viral seeds or vaccine harvests are tested, the tests are performed either at the indicated volumes or dose-equivalents. When viral seeds and vaccine harvests are tested, it is often necessary to neutralize the virus prior to inoculation. If the vaccine virus needs to be neutralized in order to perform a valid test, the source of the antiserum and its influence on test results should be considered. In general, antiserum used for neutralization should not be of human or simian origin or from the same cell type used for production of the vaccine virus.

The most appropriate tests for adventitious agents in a given vaccine will vary depending on a variety of factors, including the origin of the cell substrate and its history. We encourage manufacturers to consult with us in selecting the appropriate tests. See Section III for additional information on factors to take into account in selecting appropriate tests.

For each of the tests for adventitious agent described in this section, alternatives such as those recommended by the World Health Organization (WHO) or the European Pharmacopoeia (EP) or different inoculation strategies for testing cell bank lysates and viral harvests will be considered when justified in the context of the entire testing program.

### 1. In Vivo Tests

In vivo adventitious agent testing for vaccines includes inoculation of adult and suckling mice and inoculation of embryonated chicken eggs. Animal antibody-production tests may be performed when the potential for exposure to rodent viruses exists (e.g., through exposure to rodents or rodent cells). These tests also may be performed in other species to detect species-specific viruses. For example, inoculation of rabbits or guinea pigs may also be used to detect additional agents, when appropriate. The described animal studies must be performed in accordance with Good Laboratory Practices (GLP) regulations (21 CFR Part 58).

#### a. Adult Mice

This test detects adventitious viruses including lymphocytic choriomeningitis virus (LCMV), coxsackieviruses, flaviviruses, and rabies virus.

## Contains Nonbinding Recommendations

Each of at least 20 adult mice weighing 15-20 grams should be inoculated intraperitoneally with 0.5 mL and intracerebrally with 0.03 mL of the material to be tested. The mice should be observed daily for 21 days. Each mouse that dies after the first 24 hours of the test, or is sacrificed because of illness, should be necropsied and examined for evidence of viral infection by gross observation and by intraperitoneal and intracerebral inoculation of appropriate homogenized tissue into at least five additional mice. Each mouse inoculated with the homogenized tissue should then be observed daily for 21 days. The material may be used only if at least 80% of the originally inoculated mice and at least 80% of each subsequently inoculated group of mice remain healthy and survive the observation period, and if none of the mice show evidence of a transmissible agent or other viral infection, other than agents known to be a component of the tested material (i.e., vaccine strains of virus, when relevant) (Ref. 12).

### b. Suckling Mice

This test detects adventitious agents including many human viruses, such as coxsackievirus types A and B (type B is also detectable in cell culture) and other picornaviruses (e.g., polioviruses and echoviruses), alphaviruses, bunyaviruses (e.g., phleboviruses and nairoviruses), arenaviruses, flaviviruses, rabies, and herpesviruses (e.g., herpes simplex virus). This test can also detect many murine agents.

Each of at least 20 suckling mice less than 24 hours old should be inoculated intraperitoneally with 0.1 mL and intracerebrally with 0.01 mL of the material to be tested. The mice should be observed daily for at least 14 days. Each mouse that dies after the first 24 hours of the test, or is sacrificed because of illness, should be necropsied and examined for evidence of viral infection by gross observation and intraperitoneal and intracerebral inoculation of appropriate tissue into at least five additional mice, which should each be observed daily for 14 days. In addition, a blind passage (via intraperitoneal and intracerebral inoculation into at least 5 additional mice) should be made of a single pool of the emulsified tissue (minus skin and viscera) of all mice surviving the original 14-day test. The material may be used only if at least 80% of the originally inoculated mice and at least 80% of each group of subsequently inoculated mice remain healthy and survive the entire observation period and if none of the mice show evidence of a transmissible agent or other viral infection, other than agents known to be a component of the tested material (i.e., vaccine strains of virus, when relevant) (Ref. 12).

### c. Guinea Pigs

This test detects *Mycobacterium tuberculosis* and adventitious viruses including paramyxoviruses (including Sendai virus), reoviruses, and filoviruses.



## Contains Nonbinding Recommendations

Each of at least 5 guinea pigs each weighing 350-450 grams should be inoculated intraperitoneally with 5 mL and intracerebrally with 0.1 mL of each material to be tested. The animals should be observed daily for at least 42 days. Each animal that dies after the first 24 hours of the test, or is sacrificed because of illness, should be necropsied. All remaining animals should be sacrificed and necropsied at the end of the observation period. The material may be used only if at least 80% of the originally inoculated animals remain healthy and survive the observation period and if none of the animals shows evidence of a transmissible agent or other viral infection, other than agents known to be a component of the tested material (i.e., vaccine strains of virus, when relevant) (Ref. 12).

In vitro methods, such as culture and PCR, are also acceptable for identifying *Mycobacterium tuberculosis* when validated.

### d. Rabbits

This test detects simian herpes B virus, and should be considered when primary monkey cells are used.

Each of at least 5 healthy rabbits each weighing 1500-2500 grams should be inoculated intradermally in multiple sites with a total of 1.0 mL of the material to be tested and subcutaneously with 2.0 mL of the material to be tested. The animals should be observed daily for at least 30 days. Each animal that dies after the first 24 hours of the test, or is sacrificed because of illness, should be necropsied. The material may be used only if at least 80% of the originally inoculated animals remain healthy and survive the observation period, and if none of the animals show evidence of a transmissible agent or other viral infection, including lesions at the site of inoculation, other than agents known to be a component of the tested material (i.e., vaccine strains of virus, when relevant) (Ref. 12).

### e. Embryonated Chicken Eggs

This test detects adventitious agents including:

- by the allantoic route: orthomyxoviruses (influenza virus) and paramyxoviruses (mumps, measles, parainfluenza viruses), alphaviruses, and vesiculoviruses; and
- by the yolk sac route: herpesviruses, poxviruses, rhabdoviruses, as well as rickettsiae, mycoplasmas, and bacteria.

A sample volume, equivalent to at least 100 doses, or 10 mL, whichever represents a greater volume, should be used in egg testing. At least 10

## Contains Nonbinding Recommendations

embryonated eggs, 10 to 11 days old, should be inoculated by the allantoic route using 0.5 mL per egg. Following incubation at 35°C for 72 hours, the allantoic fluids should be harvested, pooled, and passaged by the same route into fresh, embryonated eggs, 10 to 11 days old, using 0.5 mL per egg and incubated at 35°C for 72 hours. Both the initial pool and the passaged harvest should be tested for the presence of hemagglutinating agents with red cells from guinea pigs, humans (type O), and an avian species. The tested material passes the test if at least 80% of the embryos appear normal and there is no evidence of viral or hemagglutinating agents.

At least 10 additional embryonated eggs, 6 to 7 days old, should be inoculated by the yolk sac route using 0.5 mL per egg. Following incubation at 35°C for at least 9 days, the yolk sacs should be harvested and pooled. A 10% suspension of yolk sacs should be passaged by the same route into the yolk sacs of fresh embryonated eggs, 6 to 7 days old, using 0.5 mL of inoculum per egg and incubated at 35°C for at least 9 days. The material passes the test if at least 80% of the embryos in both the initial test and the passage appear normal and there is no evidence of viral agents (Ref. 12).

### f. Antibody Production Tests

Antibody production tests are performed in the species-appropriate model (often rodents such as hamsters, rats or mice, but sometimes chickens) when the potential exists for exposure to viruses of a specific species. The test article is inoculated into specific-pathogen-free (SPF) animals that are subsequently tested for antibodies to specific agents. Agents detected in hamster antibody production (HAP) test include: lymphocytic choriomeningitis virus (LCMV), pneumonia virus of mice (PVM), reovirus type 3 (Reo3), Sendai virus, and simian virus 5 (SV5). The agents detected in rat antibody production (RAP) test include: Hantaan virus, Kilham rat virus (KRV), LCMV, mouse adenovirus, mouse encephalomyelitis virus (Theilers, GDVII), PVM, rat coronavirus (RCV), Reo3, sialodacryoadenitis virus (SDAV), Sendai virus, and Toolan virus (HI). The agents detected in mouse antibody production (MAP) test include: ectromelia virus, mouse rotavirus (EDIM) Hantaan virus, LCMV, lactic dehydrogenase virus (LDM), minute virus of mice (MVM), mouse adenovirus (MAV), murine cytomegalovirus (MCMV), mouse encephalomyelitis virus (Theilers, GDVII), mouse hepatitis virus (MHV), PVM, polyoma virus, Reo3, Sendai virus, thymic virus, and K virus (a mouse pneumonitis virus related to polyomaviruses).

A specific in vivo test for LCMV might be warranted when specific concerns about LCMV exist. This in vivo test involves intracerebral inoculation of mice used in the MAP test with live virulent LCMV to detect interference from non-lethal strains. Animals infected during the MAP test with non-virulent strains of LCMV will survive this challenge. If you use alternative

## Contains Nonbinding Recommendations

methods, such as PCR, you should demonstrate sensitivity comparable to that of the described test.

Antibody tests such as “Test for Extraneous Agents Using Chicks” (Ref. 13), might also be useful in certain cases where the vaccine virus interferes with in vitro culture assays for adventitious virus testing.

### 2. In Vitro Tests for Viruses

#### a. Cell Culture Safety Tests

Cell culture safety tests can detect a variety of adventitious viruses that include cytopathic viruses, hemadsorbing viruses, and hemagglutinating viruses. You should select cells appropriate for such tests and relevant to potential exposures of your materials including:

- monolayer cultures of the same species and tissue type as that used for production;
- monolayer cultures of human diploid cells; and
- monolayer cultures of monkey kidney cells.

Human diploid cells and monkey kidney cells are commonly used and are often susceptible to infection by many of the viruses listed below. Specialized cell culture testing might be needed to detect growth of other viruses. For example, in vitro adventitious agent testing of insect cells should be capable of detecting different families of insect viruses that could infect the host from which the cell substrate was derived. This should also include a cell line broadly susceptible to arboviruses, such as BHK21.

- Human Diploid Cells

Human viruses: adenoviruses, alphaviruses, coronaviruses, enteroviruses (including poliovirus, echovirus, and coxsackie B virus), flaviviruses, certain strains of hepatitis A virus, herpesviruses (including herpes simplex virus, varicella zoster virus, and cytomegalovirus), orthomyxoviruses, paramyxoviruses, rabies virus, reoviruses, some rhinoviruses, and rubella virus.

Simian viruses: SCMV (simian cytomegalovirus).

- Monkey Kidney Cell Line (usually Vero cells)

Human viruses: adenoviruses (some strains), alphaviruses, arenaviruses, bunyaviruses (including phleboviruses and nairoviruses), enteroviruses (including polioviruses, coxsackie B viruses, echoviruses), filoviruses, flaviviruses, herpes simplex

## Contains Nonbinding Recommendations

virus, influenza viruses, orthomyxoviruses, paramyxoviruses, polyomaviruses, poxviruses, rabies viruses, reoviruses, rhinoviruses, rotaviruses, rubella virus, and vesiculoviruses.

Simian viruses: herpes B virus.

An appropriate volume should be inoculated onto monolayer cultures of at least 3 cell types. The sample to be tested should be diluted as little as possible. The cell cultures should be observed for at least 2 weeks. After 2 weeks of observation, the original cultures of supernatants or lysates from cell banks are subcultured onto fresh cells and observed for at least an additional 2 weeks. This subculture onto fresh cells might help to distinguish between non-specific CPE and virus-induced CPE, as toxic effects of the initial specimen or length in culture will be diluted, whereas virus-induced CPE should be amplified. If the possibility of contamination with human or simian cytomegaloviruses is a consideration, the first set of human cell cultures should be observed for at least 4 weeks unless a shorter incubation period is shown to be equally sensitive. A test is considered satisfactory only if there is no evidence of adventitious agents and at least 80% of the cultures are available for observation at the end of the observation period.

The volume to be tested should be determined based on your total culture or batch size and the number of cells you require to produce a dose of vaccine. Generally, you should use cell lysate equivalent to  $10^7$  cells when testing the cell substrate. When it is not possible to directly test cells or cell lysates, a corresponding quantity of cell culture supernatant should be tested. Historically, 500 doses or 50 mL of neutralized vaccine harvest, pool, or bulk (whichever represented more virus) were used as the inoculum onto each indicator cell culture (Ref. 12). Under some circumstances, where neutralizing the vaccine strain has been difficult, smaller amounts were tested. When virus seeds are tested, it might also be necessary to neutralize the vaccine strain prior to inoculation onto indicator cells.

You may assess virus replication in cell-culture tests by various means, including development of cytopathic effect. If appropriate for your product, testing for hemadsorption or hemagglutination at the end of the observation period could facilitate the detection of non-cytopathic viruses. Hemadsorption refers to the ability of infected cells to bind to red blood cells (RBCs), while hemagglutination refers to the ability of infected cell supernatants to agglutinate RBCs. The test for hemadsorbing and hemagglutinating viruses is generally performed at the end of the observation period using guinea pig, chicken, and human type O or monkey RBCs, which are added independently to different portions of the cell cultures or to aliquots of the supernatants at 2-8°C for 30 minutes, and observed for hemadsorption and hemagglutination, respectively. This is followed by incubation with RBCs for 30 minutes at room temperature, followed by observation for

## Contains Nonbinding Recommendations

hemadsorption/hemagglutination. Red blood cells from rhesus monkeys have occasionally been used for detection of certain respiratory viruses.

The test for hemadsorbing and hemagglutinating viruses may also be performed at the end of production, directly on cells used for production or on control cell cultures (see Section III.A.4). Historically, one quarter to one third of the control vessels were tested for hemadsorbing viruses.

### b. Transmission Electron Microscopy (TEM)

TEM can detect viral particles in a cell substrate, including those from endogenous retroviruses. Under some circumstances, it might be appropriate to pre-treat cells with chemical or inducing agents to activate production of endogenous or latent viruses (Ref. 14). While TEM is fairly insensitive, it is a test that can detect adventitious agents of many types. TEM can also be used to estimate the concentration of viral particles to support validation of viral clearance. Methods include negative staining and thin section. A discussion of these methods is provided in Bierley et al. (Ref. 15).

### c. Biochemical Tests for Retroviruses

Retrovirus testing using reverse transcriptase (RT) assays should be performed on cell-free culture medium to detect retroviruses. RT assays can detect any retrovirus, as all retroviruses encode and contain RT. You should test cell substrates, viral seeds, and/or harvests of all viral vaccines produced in mammalian and insect cell substrates (or that use or include mammalian-derived materials) using a highly sensitive PCR-based RT (PBRT) assay. In certain situations, it may be appropriate to pre-treat the cells with chemical inducers of endogenous retroviruses. If you plan to use conventional RT assays, you should provide a strong justification for using a less sensitive assay.

You should validate the performance of the PBRT assay for your product, especially with regard to the lower limit of detection for both manganese- and magnesium-dependent retroviral RTs, the specificity, and the reproducibility of the assay. The lower limit of detection should be comparable with the published literature (see, for example, Refs. 16-19). If you obtain a positive result with the assay, you might need to perform infectivity studies to demonstrate whether the source of the positive result is an infectious retrovirus (Ref. 20). CBER can provide you with further information on infectivity studies upon request.

The stage of manufacture at which you should perform the PBRT test is dependent on the manufacturing process of your product. For example, products manufactured from primary cells, which are less amenable to characterization than banked cells, might need to be assessed lot-by-lot. In

## Contains Nonbinding Recommendations

cases when control cell cultures are set up in parallel with production cells (Section III.A.4), PBRT testing of cell-free medium from the control cell culture should be performed. Products manufactured from an established PBRT-tested cell bank and PBRT-tested viral seed generally do not need to be tested lot-by-lot.

CBER recognizes that some products and reagents have RT activity that does not represent adventitious infectious retroviruses (e.g., some avian cell substrates, such as chick embryo fibroblasts, chick embryo dermal cells, and eggs). For avian cells that have RT activity, you should perform testing to assure the absence of exogenous retroviruses, such as avian leukosis virus (ALV), avian sarcoma virus, and reticuloendotheliosis virus, and infectious endogenous retroviruses, such as RAV-0. You may perform this testing directly on the cell substrate. In some cases, careful documentation of the health of a specific pathogen free flock might be adequate to assure the absence of these viruses.

### d. Infectivity Tests for Retroviruses

For non-murine retroviruses, infectivity testing on indicator cells selected for their susceptibility to different retrovirus types might be appropriate. Such testing strategies should be determined on a case-by-case basis and should be discussed with CBER before implementation. You should test rodent cell substrates and viral seeds that are potentially contaminated with murine retroviruses for infectivity using *Mus dunni* cells or other highly susceptible cells and/or in S<sup>+</sup>L<sup>-</sup> assays (Ref. 21). S<sup>+</sup>L<sup>-</sup> assays may be done on feline PG4 cells (to detect amphotropic viruses), mink cells (to detect xenotropic viruses) or murine D56 cells (to detect ecotropic viruses). You may use a broadly reactive monoclonal antibody to detect viruses cultured in these assays. Infectivity tests that are capable of detecting ALV (Ref. 20) and other avian retroviruses might be appropriate for cell substrates of avian origin. Often, infectivity tests may be augmented with sensitive PBRT assays. Under some circumstances, for example when tumorigenic cell substrates are proposed for use, it might be appropriate for you to pre-treat cells with chemical agents known to induce reactivation or replication of endogenous or latent viruses (Ref. 14).

### e. PCR or Other Specific In Vitro Tests

In cases where viruses cannot be readily grown in culture, PCR is an effective tool to assess a cell substrate for contamination with such viruses. Due to the specificity of PCR, you might need to perform multiple PCR assays in order to be able to detect the full range of viruses detectable in a single more general biological assay. You should consider the tissue source and medical history of the donor from which the cell line was derived in determining what testing is appropriate. Specific PCR tests for human viruses that you should consider

## Contains Nonbinding Recommendations

include assays for hepatitis A, B, and C viruses, enteroviruses, human HIV-1, HIV-2, HTLV-1, HTLV-2, circoviruses, parvovirus B19, papillomaviruses, human polyomaviruses, human adenoviruses, Epstein-Barr virus, human cytomegalovirus, and human herpes viruses 6, 7, and 8. Specific tests for simian viruses that could potentially infect humans, including simian polyomaviruses (e.g., SV40), simian foamy virus (SFV), simian immunodeficiency virus (SIV), simian retrovirus (SRV), and simian T-cell lymphotropic virus (STLV), should also be considered. You also should consider using PCR assays that detect several agents using degenerate or consensus primers provided that the sensitivity of these assays is sufficient to improve assurance of product safety. PCR testing also can play an important role in qualification of insect cell substrates.

Although less sensitive, you also may consider the use of Southern blot hybridization analysis of DNA extracted from the cell substrate or tissue culture to assess for contamination with some viruses. Either PCR or Southern hybridization may be used to detect non-cytopathic agents in tissue culture tests.

### f. Sourcing and Testing of Bovine-Derived Materials

You should source bovine-derived reagents in a manner that minimizes the risk of contamination from BSE (Ref. 33). We recommend that you consult with CBER for more information. In some cases, a BSE risk assessment can address BSE-related issues.

We recommend that you test bovine-derived reagents using assays at least as sensitive as the cell culture and fluorescent antibody testing recommended in 9 CFR 113.53 (Requirements for ingredients of animal origin used for production of biologics) and 113.47 (Detection of extraneous viruses by the fluorescent antibody technique). We recommend the use of Vero cells and bovine cell culture, such as bovine turbinate cells. Agents of concern from bovine sources include: bovine parainfluenza Type 3, infectious bovine rhinotracheitis virus, rabies virus, reovirus, bovine adenoviruses, bovine respiratory syncytial virus, bovine parvovirus, bluetongue virus, and bovine viral diarrhea virus (including non-cytopathic strains).

We may request testing for other potential contaminants of bovine materials depending upon exposure history and species of origin of a cell substrate or reagent including circoviruses, bovine herpesviruses, bovine polyomavirus, bunyaviruses, and bovine retroviruses (such as bovine immunodeficiency virus and bovine leukemia virus). Some of these agents might be detected in cell culture, while others might require additional tests.

## Contains Nonbinding Recommendations

### g. Testing of Porcine-Derived Reagents

Products that will be administered to humans should not be contaminated with animal adventitious agents. Products should thus be free of the agents listed in the requirements for porcine adventitious virus testing of products intended for use in animals (9 CFR 113.47). Testing of porcine-derived reagents for these agents is an important component of providing this assurance. Under certain circumstances, additional tests might be needed. Agents of concern from porcine cells listed in 9 CFR 113.47(b)(1) and (6) include: porcine parvovirus, bovine viral diarrhea virus (BVDV), reovirus, rabies virus, porcine adenovirus, transmissible gastroenteritis virus (TGE), and porcine hemagglutinating encephalitis virus. At a minimum, you should test porcine trypsin for porcine parvovirus (PPV).

Other porcine agents that might be of concern, depending upon exposure history and species of origin of your cell substrate or reagent, include circoviruses, enteroviruses, porcine respiratory and reproductive syndrome virus (PRRS), porcine cytomegalovirus, porcine influenza viruses, pseudorabies virus, swine pox virus, swine fever virus (hog cholera virus, African), vesicular stomatitis virus (VSV) (Ref. 22), Nipah virus and porcine retroviruses.

### 3. In Vitro Tests for Non-Viral Agents

#### a. Mycoplasma/Spiroplasma

All licensed viral vaccines produced in cell substrates (and similar products) should be tested to ensure the absence of mycoplasma contamination. Testing is usually performed on the virus seed and/or MCBs, cell substrate, and a representative portion of each working cell stock used for manufacture of the product. Each lot of product harvest concentrate should be sampled for testing prior to further processing, *e.g.*, prior to clarification, filtration, purification, and inactivation, unless testing at this stage of the manufacturing process is not feasible. Under 21 CFR 610.30, for virus vaccines produced from in vitro living cell cultures, you must test each virus harvest pool and control fluid pool (when control cells are used) for mycoplasma prior to clarification or filtration in the case of live virus vaccines and prior to inactivation in the case of inactivated virus vaccines. Prior to testing, the product harvest concentrate sample must be stored either: 1) between 2 and 8°C for no longer than 24 hours, or 2) at -20°C or lower if stored for more than 24 hours. If stored for longer than 24 hours, we recommend that you store the sample at -60°C or lower to allow for greater recovery of mycoplasma.

Although 21 CFR 610.30 specifies agar and broth media culture procedures, recent data suggest that some fastidious strains of mycoplasma, especially *Mycoplasma hyorhinis*, might not be detected by these agar and broth media



## Contains Nonbinding Recommendations

procedures. Therefore, you should include an indicator cell culture procedure to detect such strains. We consider the testing described below to be acceptable alternatives to the method specified in 21 CFR 610.30. However, you must seek and obtain FDA approval for these modifications in accordance with 21 CFR 610.9(b). Additional information regarding mycoplasma testing is available (Ref. 23).

For Spiroplasma testing by the culture method and by DNA staining, you should use media that supports the growth of Spiroplasma, Entomoplasma, and Mesoplasma.

### Testing for Mycoplasma using both Alternative Agar and Broth Media, and Indicator Cell Culture Procedures

- Agar and Broth Media Procedure

(1) Each lot of agar and broth medium should be free of antibiotics except for penicillin, and each lot of medium should be examined for mycoplasma growth-promoting properties. To demonstrate the capability of the media to detect known mycoplasma contaminants, use the mycoplasma cultures specified below in (3)(i) as positive controls.

(2) (i) Inoculate no less than 0.2 mL of the test sample (e.g., product harvest concentrate sample) in evenly distributed amounts over the surface of 2 or more agar plates of 1 medium formulation.

(ii) Inoculate no less than 10 mL of the product harvest concentrate sample into a flask containing 50 ml of broth medium which is incubated at  $36 \pm 1^\circ \text{C}$ .

(iii) Test 0.2 mL of the broth culture on the 3rd, 7th, and 14th days of incubation by subculture onto 2 or more agar plates of the same medium formulation as that used above in (i).

(iv) Incubate 2 of the initial isolation plates and 2 each of the three subculture plates in a 5 to 10 % carbon dioxide in nitrogen and/or hydrogen atmosphere containing less than 0.5% oxygen during the test incubation period.

(v) Incubate all culture agar plates for no fewer than 14 days at  $36 \pm 1^\circ \text{C}$  and observe them microscopically at 100 times magnification or greater for growth of mycoplasma colonies.

(3) (i) Include in each test at least 2 known mycoplasma species or strains as positive controls, 1 of which should be a dextrose

## Contains Nonbinding Recommendations

fermenter (i.e., *M. pneumoniae* strain FH or equivalent species or strains) and 1 of which should be an arginine hydrolyzer (i.e., *M. orale* strain CH19299 or equivalent species or strain). Positive control cultures should be not more than 15 passages from isolation and should be used in a standard inoculum of 100 colony-forming units (CFU) or 100 color-changing units (CCU) or less.

(ii) Include uninoculated agar medium as a negative control.

(4) Interpret the results of the procedure according to the specification detailed in this section under Interpretation of Results.

- Indicator Cell Culture Procedure

(1) Using a Vero cell culture substrate, pre-test the procedure by using the mycoplasma cultures specified below in (3)(i) as positive controls to demonstrate the capability of the cell substrate to detect known fastidious mycoplasmal contaminants. An equivalent indicator cell substrate might be acceptable if data demonstrate at least equal sensitivity for the detection of known mycoplasmal contaminants.

(2) (i) Inoculate no less than 1 mL of the test sample (e.g., product harvest concentrate sample) to 2 or more indicator cell cultures grown on cover slips in dishes or equivalent containers.

(ii) Incubate the cell cultures for 3 to 5 days at  $36 \pm 1^\circ\text{C}$  in a 5% carbon dioxide atmosphere. Examine the cell cultures for the presence of mycoplasmas by epifluorescence microscopy using a DNA-binding fluorochrome, such as bisbenzimidazole or an equivalent stain.

(3) (i) Include in each test two known mycoplasma species or strains as positive controls (i.e., *M. hyorhinis* strain DBS 1050, *M. orale* strain CH19299, or equivalent species and strains), using an inoculum of 100 CFU or 100 CCU or fewer.

(ii) Include as a negative control a non-infected indicator cell culture.

(4) Interpret the results of the procedure according to the specifications detailed in this section under Interpretation of Results.

## Contains Nonbinding Recommendations

- Interpretation of Results

(1) For the agar and broth media procedure, compare the appearance of the media inoculated with the product to that of the positive and negative controls.

(2) For the indicator cell culture procedure, using 600 times magnifications or greater, compare the microscopic appearance of the cultures inoculated with the product to that of the positive and negative cell controls.

(3) Marked cytopathic effects or nuclear chromatin fragmentation caused by vaccine virus infection that affect the interpretation of the results can be minimized by using a specific neutralizing viral antiserum or a nonpermissive cell culture substrate. The antisera should also be added to the positive and negative controls.

(4) The product is considered satisfactory for manufacture if both the agar and/or broth media procedure and the indicator cell culture procedure show no evidence of mycoplasma contamination (*i.e.*, no growth) and thus resemble the negative control(s) for agar and broth and indicator cell procedures.

(5) If mycoplasmas are recovered, you should perform confirmatory testing to establish the species.

### Testing for Mycoplasma Using PCR-based Assays

PCR-based assays may be used to detect mycoplasma, provided that such an assay can be shown to be comparable to the agar and broth media procedure and the indicator cell culture procedure. Such assays may combine culture and PCR. In some cases, culture-based procedures cannot be used due to an inability to completely neutralize vaccine viruses, thus necessitating the use of PCR-based assays to test for mycoplasma in these products.

#### b. Bacterial and Fungal Sterility

You must perform the test for bacterial and fungal sterility described in 21 CFR 610.12 or an equivalent method that fulfills the requirements of 21 CFR 610.9.

#### c. Mycobacteria Testing

A 2.0 mL sample of the filtrate in its most concentrated form should be inoculated onto Lowenstein – Jensen’s egg medium or other media

## Contains Nonbinding Recommendations

demonstrated to be equally capable of supporting growth. An appropriate positive-control test should be conducted simultaneously with the sample under test, and the test should be shown to be capable of supporting the growth of small numbers of the mycobacteria. All the test vessels should be incubated at a suitable temperature for a period of 6 weeks under conditions that will prevent drying of the medium, after which the cultures should be examined for evidence of mycobacterial colonies. The filtrate is satisfactory for use if the test shows no evidence of mycobacteria (Ref. 12).

Guinea pig tests have also been used to detect mycobacteria. If you use alternative tests, including PCR assays, they should be shown to be sufficiently sensitive to test for the presence of mycobacteria.

### **B. TESTING OF CELL PROPERTIES**

Tests for tumorigenicity and oncogenicity as outlined in the section below are required to follow good laboratory practices for conducting nonclinical laboratory studies as described in the GLP regulations (21 CFR Part 58).

#### 1. Tests for Tumorigenicity

Tumorigenicity is defined as the process by which cells form tumors when inoculated into animals (generally a syngeneic, an immunosuppressed allogeneic or an immunosuppressed xenogeneic host) (see Section V. Glossary). Tumorigenicity is a characteristic of the immortalized cells themselves, rather than of the agents or components present in them. We recommend you contact CBER for advice as to the need for tumorigenicity testing.

Tumorigenic cells have not traditionally been used for the production of prophylactic viral vaccines, primarily because of theoretical concerns that components within tumorigenic cells could induce tumors in vaccine recipients. These concerns include the potential presence of exogenous agents, such as oncogenic viruses, and the potential risk from endogenous materials, such as endogenous viruses or oncogenic nucleic acids. In addition, intact human cells derived from human tumors have been shown to form tumors in allogeneic humans.

The goal in tumorigenicity testing is to determine whether your cell substrate is capable of forming tumors after inoculation into animals. The TPD<sub>50</sub> (tumor-producing dose in 50% of animals) and capacity to form metastases are characteristic properties of a cell line, and these characteristics might be used to further define the tumorigenic phenotype of a cell line. Considerations associated with tumorigenicity testing include: 1) choice of appropriate animal models; 2) definition of a positive result; 3) determination of the appropriate duration of testing; 4) determination of appropriate numbers of cells to be tested; and 5) selection of appropriate controls.

## Contains Nonbinding Recommendations

You should use an animal model that is known to be susceptible to tumor formation by tumorigenic cells. Because immunocompromised adult and newborn rodents are relatively sensitive for revealing a tumorigenic phenotype, you should consider these animal models. Thus, the most commonly used animals for tumorigenicity testing are nude (nu/nu) mice because they are T-cell deficient. Newborn nude mice appear to be more susceptible to tumor formation than adult nude mice (Ref. 24), suggesting that newborn nude mice might be the best choice to use when identification of a weakly tumorigenic phenotype is important. You might choose to use another animal model if it has been shown to have comparable sensitivity to the nude mouse model.

Regardless of the test system you use, the animals should be observed and palpated at regular and frequent intervals for the formation of nodules at the site of injection. Any nodules formed should be measured in two dimensions. We consider progressive tumor formation at the site of injection a positive result. Some cell types might cause tumors at distant sites as well (Ref. 24), and these also should be reported. Animals with nodules that begin to regress during the period of observation should not be sacrificed, as nodules that spontaneously regress do not represent progressively growing tumors and are not indicative of a tumorigenic phenotype. You may euthanize any animals with tumors before the end of the study if the tumor interferes with the comfort of the animal. Intercurrent infections, including subclinical infections, have the potential to alter the incidence of tumors, affecting the quality of the data and interfering with the purpose of the study (Ref. 25).

You should perform a necropsy on each animal when it dies, when it is euthanized, or at the end of the observation period. The necropsy should include examination for gross evidence of tumor formation at the site of inoculation and in all major organ systems such as lymph nodes, lungs, brain, spleen, kidneys, and liver. All lesions, detectable regional lymph nodes, the site of inoculation, and the lungs are to be examined histologically. In some cases (for example, tumors distant from the site of inoculation), it might be necessary for you to use molecular or immunological methods to distinguish cell-substrate-related from spontaneous tumors.

The inoculum per animal should consist of  $10^7$  test cells or positive control tumor cells suspended in 0.2 mL (0.1 mL for newborn animals) volume of serum-free medium administered by the subcutaneous route. You should inoculate at least ten animals, each with  $10^7$  test cells that are at or beyond the end-of-production passage level and at least ten animals with your positive control tumor cells. At least 9 out of 10 animals injected with positive control cells (e.g., HeLa cells or other cells with comparable tumorigenicity) should show progressively growing tumors in order for your test to be valid.

## Contains Nonbinding Recommendations

Selection of the appropriate duration of testing requires that you balance the increased sensitivity that might be obtained using a longer test, against the likelihood of false-positive results due to spontaneous tumor formation. Weakly tumorigenic cells might require between 4 and 7 months to form tumors in nude mice. Thus, extended observation periods might be necessary in some cases. CBER can provide you with further information on tumorigenicity testing and the appropriate observation periods.

### 2. Testing for Oncogenicity

Oncogenicity testing (see Section V. Glossary) is designed to assure that agents that could immortalize cells and endow them with the capacity to form tumors are not present in a cell substrate. If your vaccine is manufactured in a cell substrate that was derived from a tumor, or that has a tumorigenic phenotype through an unknown mechanism, it might carry a higher theoretical risk of containing oncogenic substances.

If the presence of an oncogenic virus is suspected because of the cell phenotype or the origin of your cell substrate, we recommend that you perform oncogenicity testing in animals using lysates of the cell substrate. For cell substrates with a tumorigenic phenotype, we recommend that you perform oncogenicity testing in animals using DNA from the cell substrate in order to provide assurance that residual DNA is non-oncogenic (also see Section III.C.6). Oncogenicity testing might also be appropriate for some products with high quantities of residual cellular DNA. Existing assays for the presence of oncogenic agents in certain cell substrates might not be adequate to provide sufficient assurance of the safety of your vaccine for clinical use. You should consult with CBER if you have any questions on the necessity of these tests and the methods to be followed.

All tumors that develop in oncogenicity assays should be examined for the presence of DNA from the species that the cell substrate was derived from, and of the animal in which oncogenicity testing was performed. We recommend retention of samples from any tumors observed in oncogenicity assays to permit further investigation when necessary.

### 3. Identity Testing of Cell Substrates

Identity testing is required by the general requirements for laboratory controls (21 CFR 211.160(b)) and the general standards for biological products (21 CFR 610.18(b)). Identity tests assure that a licensed viral vaccine is as stated on the manufacturer's label. Identity tests also provide extra assurance in facilities or laboratories that are propagating multiple cell lines or viruses.

Biochemical tests such as polyacrylamide gel electrophoresis (PAGE) analysis may be used to determine the species from which the cell substrate was derived. Comparison of the migration patterns of the cellular enzymes malate

## **Contains Nonbinding Recommendations**

dehydrogenase, glucose-6-phosphate dehydrogenase, nucleoside phosphorylase and lactate dehydrogenase is another method (Ref. 26).

Cytogenetic analysis can be used to determine the karyotype of the cell substrate as a means of identifying the species, sex, ploidy, and, grossly, genetic stability. You should examine 100-200 metaphase cells. However, it might be appropriate for you to examine more than 200 such cells, depending on the heterogeneity of the karyotypes of your product.

DNA fingerprinting is the most commonly used test of genetic identity and has been used to document cell origin and to determine whether cell contamination has occurred (Ref. 27). Specific methods for this purpose can be found in Stacey, et al. (Ref. 28).

Tests such as Southern blot hybridization, PCR, growth under selective conditions, expression of a reporter gene or other gene of interest, etc., may be applied to distinguish an engineered cell line both from the parent cell line and from other cell lines. Some examples can be found in Stacey, et al., (Ref. 29). You may also use other tests that provide information relevant to identity of cells, such as HLA typing.

### **4. Testing for Genetic Stability**

You should demonstrate the genetic stability of your cell substrate from the establishment of the MCB through and perhaps beyond the end of production. For an engineered cell line, the inserted gene of interest should remain intact and at the same copy number, and be expressed at comparable levels throughout production. Also, a diploid cell strain should remain diploid throughout. If such characteristics are not stable, then you should demonstrate that the instability does not adversely impact manufacturing or product consistency. For methods to assess a cell substrate's genetic stability, reference is made to the ICH Q5B and Q5D documents (Refs. 30 and 3, respectively).

## **C. OTHER TESTS**

### **1. Testing for the Presence of Residual Cells**

You should assure that your final vaccine product does not contain residual cells. Processes, such as filtration, should be implemented and validated to ensure that intact cells are not present in the final product. Validation that residual cell removal processes are robust is important for immortalized cells. Determining the extent to which intact cells (or other materials known to be smaller than intact cells) are cleared by these processes is an important part of this validation.

## Contains Nonbinding Recommendations

### 2. Testing for Residual Cellular DNA

Residual DNA might be a risk to your final product because of oncogenic and/or infectivity potential. There are several potential mechanisms by which residual DNA could be oncogenic, including the integration and expression of encoded oncogenes or insertional mutagenesis following DNA integration. Residual DNA also might be capable of transmitting viral infections if retroviral proviruses, integrated copies of DNA viruses, or extrachromosomal genomes are present.

The risks of oncogenicity and infectivity of your cell-substrate DNA can be lessened by decreasing its biological activity. This can be accomplished by decreasing the amount of residual DNA and reducing the size of the DNA (e.g., by DNase treatment or other methods) to below the size of a functional gene (based on current evidence, approximately 200 base pairs). Chemical inactivation can decrease both the size and biological activity of DNA. If DNA removal, digestion, or inactivation is undertaken, you should validate your methods.

You should measure the amount and size distribution of residual DNA in your final product. For widely used human diploid cell strains, such as MRC-5 and WI-38 cells, measurement of residual DNA might be unnecessary because we do not consider residual DNA from these human diploid cells to be a safety issue. We might require limitation of the amount of residual DNA, depending on the potential risks associated with that DNA, for human diploid or primary cell types for which there is less experience. You should limit residual DNA for continuous non-tumorigenic cells, such as low-passage Vero cells, to less than 10 ng/dose for parenteral inoculation as recommended by WHO (Ref. 31). Because orally administered DNA is taken up approximately 10,000-fold less efficiently than parenterally administered DNA, we recommend limiting DNA to less than 100 µg/dose for oral vaccines (Ref. 32). If you are using cells with tumorigenic phenotypes or other characteristics that give rise to special concerns, more stringent limitation of residual DNA quantities might be needed to assure product safety.

### 3. General Safety Test (GST)

The requirements for, and the exceptions and criteria for exception from, the GST are set forth in 21 CFR 610.11.



## Contains Nonbinding Recommendations

### V. GLOSSARY

Because different guidance documents might use the same or similar terms differently, the following glossary is provided to define the terminology used in this document.

1. **ADVENTITIOUS AGENT:** A microorganism (including bacteria, fungi, mycoplasma/spiroplasma, mycobacteria, rickettsia, viruses, protozoa, parasites, TSE agent) that is inadvertently introduced into the production of a biological product.
2. **ANCILLARY REAGENT:** Reagents that are used in the manufacture or production of a biological product that may or may not end up as part of the final product. Examples include: insulin, transferrin, growth factors, interferon, interleukins, and other proteins, drugs or chemicals like dimethyl sulfoxide.
3. **ANEUPLOID:** Having an atypical number of chromosomes which is not an integer multiple of the haploid number.
4. **CELL BANK:** Vials of cells of uniform composition (although not necessarily clonal) derived from a single tissue or cell, aliquoted into appropriate storage containers, and stored under appropriate conditions, such as the vapor phase of liquid nitrogen.
5. **CELL LINE (CL):** Cells that have been propagated in culture since establishment of a primary culture and survival through crisis and senescence. Such surviving cells are immortal and will not senesce. Diploid cell strains have been established from primary cultures and expanded into cell banks but have not passed through crisis and are not immortal. [The ATCC uses the abbreviation CCL to signify their Certified Cell Lines.]
6. **CHARACTERIZATION:** Determination of the properties of a cell substrate or cell bank.
7. **CONTROL CELLS:** Cells that are split off from the production culture and maintained in parallel under the same conditions and using the same reagents (e.g., culture medium) in order to perform tests on cells that have not been exposed to the vaccine virus (which may interfere with some tests).
8. **DIPLOID:** Having the expected number of chromosomes for a species, (i.e., two of each autosomal chromosome and two sex chromosomes).
9. **ENDOGENOUS VIRUS:** A virus whose genome is present in an integrated form in a cell substrate by heredity. Endogenous viral sequences may or may not encode an intact or infectious virus.
10. **END-OF-PRODUCTION CELLS (EOPC):** Cells cultured (under conditions comparable to those used in production) from the MCB or WCB to a passage level or population doubling level comparable to or beyond the highest level reached in production.

## Contains Nonbinding Recommendations

11. **FINAL BULK:** The stage of vaccine production directly prior to filling of individual vials. The final bulk often represents a processed harvest.
12. **FREE OF and FREEDOM FROM:** For a substance to be considered free of a contaminant, an assay must demonstrate that a defined quantity of the substance is negative for that contaminant to a defined level of sensitivity. The level of assay sensitivity is defined by the choice of assay and can be determined experimentally using standardized reagents. Alternatively, a validated process that is known to remove a contaminant to a defined level may be used to demonstrate freedom from that contaminant.
13. **HARVEST:** At the end of vaccine virus propagation in cell culture, material is collected from which vaccine will be prepared. This material may be the culture supernatant, the cells themselves (often in disrupted form), or some combination thereof.
14. **IMMORTALIZATION:** The process by which cells with finite lifespan (e.g., primary cells, diploid cell strains) are converted to those with infinite lifespan.
15. **LATENT VIRUS:** A virus that is present in a cell, without evidence of active replication, but with the potential to reactivate, is considered to be microbiologically latent.
16. **MANUFACTURER'S WORKING CELL BANK (MWCB) OR WORKING CELL BANK (WCB):** A cell bank derived by propagation of cells from MCB under defined conditions and used to initiate production cell cultures on a lot-by-lot basis.
17. **MASTER CELL BANK (MCB):** A bank of a cell substrate from which all subsequent cell banks used for vaccine production will be derived. The MCB represents a characterized collection of cells derived from a single tissue or cell.
18. **MASTER VIRUS SEED (MVS):** A viral seed of a selected vaccine virus from which all future vaccine production will be derived, either directly, or via Working Virus Seeds.
19. **ONCOGENICITY:** The property of certain biological agents (e.g., viruses) or materials (e.g., nucleic acids) that are capable of immortalizing cells and endowing them with the capacity to form tumors. Oncogenicity is distinct from tumorigenicity (See Tumorigenicity).
20. **PARENTAL VIRUS:** A virus that has been manipulated in some manner to generate a viral seed with characteristics needed for production.
21. **PARENT CELL BANK:** A few vials consisting of cells from which the Master Cell Bank was derived. Parental Cells may be manipulated to derive a cell substrate with desired characteristics.
22. **PASSAGE LEVEL:** The number of times, since establishment from a primary cell culture, a culture has been split or re-seeded.

## Contains Nonbinding Recommendations

23. **POPULATION DOUBLING LEVEL:** The number of times, since establishment from a defined point in the history of a cell substrate (often the primary cell culture), a culture has doubled in number of cells.
24. **PRIMARY CELLS:** Cells placed into culture immediately after an embryo, tissue, or organ is removed from an animal or human and homogenized, minced, or otherwise separated into a suspension of cells.
25. **PURITY:** Relative freedom from extraneous matter in the finished product, whether or not harmful to the recipient or deleterious to the product. (21 CFR 600.3(r))
26. **QUALIFICATION:** Determination of the suitability of a cell substrate for manufacturing based on its characterization.
27. **TUMORIGENIC:** A cell type is tumorigenic if it forms tumors when inoculated into animals (generally a syngeneic, an immunosuppressed allogeneic, or an immunosuppressed xenogeneic host). These tumors may be at the injection site or a different site and may also metastasize to other sites.
28. **NON-TUMORIGENIC:** A cell type is non-tumorigenic if it is shown not to form tumors in appropriate animal models.
29. **TUMORIGENICITY:** Tumorigenicity is the process by which immortalized cells form tumors when inoculated into animals (see Tumorigenic). Tumorigenicity is distinct from Oncogenicity (See Oncogenicity).
30. **TUMORIGENICITY TESTING:** An assay/test that determines whether or not immortalized cells are tumorigenic when injected into animals.
31. **VALIDATION OF ANALYTICAL PROCEDURES:** Validation defines the performance characteristics of an analytical procedure, based on the demonstration that the procedure is suitable for its intended purpose or use. Validation is generally performed in accordance with the relevant ICH guidelines.
32. **VALIDATION OF PROCESSES:** Process validation is establishing documented evidence that provides a high degree of assurance that a specific process will consistently produce a product meeting its pre-determined specifications and quality characteristics.
33. **VIRAL CLEARANCE:** The combination of the physical removal of viral particles and the reduction of viral infectivity through inactivation.
34. **VIRUS PRE-SEED:** A few vials consisting of vaccine virus from which the MVS was derived.

## **Contains Nonbinding Recommendations**

35. **VIRUS SEED or VIRAL SEED:** A live viral preparation of uniform composition (although not necessarily clonal) derived from a single culture process, aliquoted into appropriate storage containers, and stored under appropriate conditions.
36. **WORKING CELL BANK (WCB):** See “Manufacturer’s Working Cell Bank (MWCB)”.
37. **WORKING VIRUS SEED (WVS):** A viral seed derived by propagation of virus from the MVS under defined conditions and used to initiate production cell cultures lot-by-lot.

## Contains Nonbinding Recommendations

### VI. REFERENCE LIST

1. Points to Consider in the Characterization of Cell Lines Used to Produce Biologicals, July 12, 1993.
2. The International Conference on Harmonisation (ICH) Guidance, *Q5A: Viral Safety Evaluation of Biotechnology Products Derived from Cell Lines of Human or Animal Origin*, (63 FR 51074; September 24, 1998).
3. ICH Guidance, *Q5D: Derivation and Characterisation of Cell Substrates Used for Production of Biotechnological/Biological Products*, (63 FR 50244; September 21, 1998).
4. Goffe, A.P., Hale J, and Gardner, P.S.: Poliomyelitis Vaccine, Letters to Editor. *Lancet* 1961; 1:612.
5. Garnick R.L., Raw Materials as a Source of Contamination in Large-Scale Cell Culture, *Dev Biol Stand.* 1998; 93:21-29.
6. The International Conference on Harmonisation (ICH) Guideline, *Q2(R1): Validation of Analytical Procedures: Text and Methodology*, (60 FR 11260, March 1, 1995) and (62 FR 27463, May 19, 1997).
7. U.S. Pharmacopeia (USP) Chapter 1225: Validation of Compendial Methods.
8. Rezapkin G.V., et al., Microevolution of Sabin 1 strain *in vitro* and genetic stability of oral poliovirus vaccine, *Virology* 1994; 202(1): 370-378.
9. Guidance for Industry: Source Animal, Product, Preclinical, and Clinical Issues Concerning the Use of Xenotransplantation Products in Humans, April 2003. (see also 21 CFR 610.18(c)(1)(iii)).
10. European Pharmacopoeia, 4th Edition, Section 2.6.6.
11. U.S. Pharmacopeia, Chapter 61: Microbial Examination of Non Sterile Products: Microbial Enumeration Tests.
12. Office of the Federal Register, Code of Federal Regulations, 21 CFR part 630 (Additional Standards for Viral Vaccines). 1996 and earlier versions.
13. European Pharmacopoeia, 4<sup>th</sup> Edition, Test for Extraneous Agents Using Chicks, Section 2.6.6.
14. Khan, A.S., Muller, J., and Sears, J.F., Early detection of endogenous retroviruses in chemically induced mouse cells, *Virus Research* 2001; 79:39-45.

## Contains Nonbinding Recommendations

15. Bierley, S.T. et al., A comparison of methods for the estimation of retroviral burden., *Dev. Biol. Stand.* 1996; 88:163-5.
16. Arnold et al., One-step fluorescent probe product-enhanced reverse transcriptase assay, *Biotechniques* 1998; 25: 98-106.
17. Maudru and Peden, Adaptation of the fluorogenic 5'-nuclease chemistry to a PCR-based reverse transcriptase assay, *Biotechniques* 1998; 25: 972-975.
18. Lovatt, et al., High throughput detection of retrovirus-associated reverse transcriptase using an improved fluorescent product enhanced reverse transcriptase assay and its comparison to conventional detection methods, *J Virol Methods* 1999; 82: 185-200.
19. Sears, J.F. and Khan, A.S., Single-tube fluorescent product-enhanced reverse transcriptase assay with Ampliwax (STF-PERT) for retrovirus quantitation, *J Virol Methods* 2003; 108(1):139-42.
20. Khan AS, et al., The reverse transcriptase activity in cell-free medium of chicken embryo fibroblast cultures is not associated with a replication competent retrovirus, *J Clin Virol* 1998; 11:7-18.
21. European Pharmacopoeia, 4<sup>th</sup> Edition, Section 2.6.4.
22. Leman, A.D., et al., *Diseases of the Swine*, 1992, 7th Edition.
23. Barile, M.F. and Rottem, S., *Mycoplasmas in Cell Culture, Rapid Diagnosis of Mycoplasmas*, I. Kahane and A. Adoni (ed.) Plenum Press, New York, 1993; p.155-193.
24. Sordat, B., et al., Invasive Growth and Dissemination of Human Solid Tumors and Malignant Cell Lines Grafted Subcutaneously to Newborn Nude Mice, *Proceedings of the Second International Workshop on Nude Mice* 1977; p. 313-326.
25. National Research Council (Commission on Life Sciences, Institute of Laboratory Animal Resources), *Surveillance, Diagnosis, Treatment and Control of Disease in Guide for the Care and Use of Laboratory Animals*, National Academy Press, Washington DC, 1996, pp. 59-60.
26. Stacey, G.N., et al., Authentication of Animal Cell Cultures by Direct Visualization of Repetitive DNA, Aldolase Gene PCR and Isoenzyme Analysis, *Biologicals* 1997; 25:75-85.
27. Stacey, G.N., et al, DNA Fingerprinting Transforms the Art of Cell Authentication, *Nature*, 1992; 357:261-262.
28. Stacey, G.N., et al., The Quality Control of Cell Banks Using DNA Fingerprinting, *EXS.* 1991; 58:361-70.

## Contains Nonbinding Recommendations

29. Stacey G.N., Phillips, P., Quality Assurance for Cell Substrates, *Dev Biol Stand.* 1999; 98:141-51.
30. ICH Guidelines, *Q5B: Quality of Biotechnological Products: Analysis of the Expression Construct in Cells Used for Production of r-DNA Derived Protein Products*, (61 FR 7006, February 23, 1996).
31. World Health Organization, *Requirements for the use of animal cells as in vitro substrates for the production of biologicals*. World Health Organization (WHO) Technical Report. Series 878, 1998, Annex 1
32. World Health Organization, *Guidelines To Assure The Quality, Safety And Efficacy Of Live Attenuated Rotavirus Vaccines (Oral)*, World Health Organization (WHO) Technical Report Series 941, 2007, Annex 3
33. Proposed Rule, *Use of Materials Derived from Cattle in Medical Products Intended for Use in Humans and Drugs Intended for Use in Ruminants*, (72 FR 2007 1582; Jan. 12, 2007).

## **Contains Nonbinding Recommendations**

### Appendix 1

Multi-stage testing with a variety of different assays is used to assure product safety and ensure product quality. This includes testing all the biological materials in vaccine manufacture, especially the initial components, which are the virus seed and the cell substrate. Different stages for testing are selected based on the maximum likelihood of adventitious agent detection and effect on product quality. This table shows a typical testing scheme; actual selected tests will depend upon the actual cell substrate, the product, and its manufacture. Additional tests may be included on a case-by-case basis depending upon the cell substrate and other biological materials as well as production conditions for vaccine manufacture.



### Contains Nonbinding Recommendations

Table 1: Example of a Biosafety Testing Scheme for Manufacture of a Viral Vaccine.

	Cell Testing				Testing of Intermediates and Final				
	MCB	WCB	Control Cells	EOP	Virus Seed Harvest	Virus Seed Final	Vaccine Harvest	Vaccine Bulk	Final Filled
IDENTITY & PURITY <sup>1</sup>	+	(+)		(+)	(+)	+		(+)	+
STABILITY (genotype/phenotype/serotype)	(+)			(+)		(+)			
STERILITY									
Bacteria and Fungal	+	+	+	+	+	+	+	+	+
Mycoplasma or Spiroplasma	+	+	+	+	+	+	+		
Mycobacteria	(+)	(+)	(+)	(+)	(+) <sup>2</sup>		(+)		
ADVENTITIOUS AGENT TESTING									
Cytopathic and Hemadsorbing viruses			+						
<i>In vitro</i> cell cultures	+	+	(+)	+	+		(+)		
<i>In vivo</i> assays	+		(+)	+	+		(+)		
TEM <sup>4</sup>	+			+					
Antibody Production assay	(+)				(+)				
Bovine viruses	(+)	(+)	(+)	(+)	(+)		(+)		
Porcine viruses	(+)	(+)	(+)	(+)	(+)		(+)		
Retroviruses <sup>4</sup>	+		(+)	+	+		(+)		
Tests for specific viruses <sup>4</sup>	(+)	(+)	(+) <sup>3</sup>		(+)		(+)		
TUMORIGENICITY				+					
ONCOGENICITY				(+)					
OTHER AGENTS									
Residual cell DNA								+	
Residual cellular protein								+	
Residual serum, antibiotics, benzoylase, ancillary reagents <sup>2</sup>								+	
GENERAL SAFETY TEST									+
ENDOTOXIN									+

## Contains Nonbinding Recommendations

<sup>1</sup> tests may be different for cells versus intermediate and final products

<sup>2</sup> if applicable

<sup>3</sup> including ALV test in case of chicken cells

<sup>4</sup> may be combined with virus induction assays, as applicable

+ required stage for testing

(+) alternative stage for testing